

4D microscopy of red blood cell membrane biophysics during *Plasmodium falciparum* invasion.

Niall D Geoghegan^{1,2}, Cindy Evelyn¹, Lachlan Whitehead^{1,2}, Michal Pasternak^{1,2}, Julie Healer^{1,2}, Alan Cowman^{1,2} and Kelly Rogers^{1,2}.

The Walter and Eliza Hall Institute of Medical Research¹
The University of Melbourne²
1G Royal Parade, Parkville, Melbourne, VIC, Australia 3052
E-mail: Geoghegan.n@wehi.edu.au

KEY WORDS: 4D Imaging, Lattice-light Sheet Microscopy, Live Cell Imaging, Membrane Biophysics

The symptoms of malaria are manifested during asexual blood stage of *Plasmodium* parasites, at the time of rapid multiplication of the parasite. This time window is a common target for the development of antimalarial therapies and potential vaccines. Difficulties in the development of efficient anti-malaria vaccines lie, in part, due to a fundamental lack of insight into the molecular processes and biophysical mechanisms which govern host-pathogen interactions at this stage. To this end invasion of the red blood cell has been studied by various microscopy techniques at fixed points in time [1, 2, 3]. The biophysical mechanisms for invasion have also been suggested through combined experimentation and modelling [4]. However, these studies lack the necessary temporal resolution to fully understand the molecular and biophysical basis for invasion in real time.

An important part of the invasion process is the formation of the parasitophorous vacuole membrane (PVM) at the point of entry to the host red blood cell. This membrane provides a physical barrier and an exchange surface between the parasite and the host cell. The formation of the PVM and subsequent remodelling of the host membrane during invasion are incredibly dynamic events and are very challenging to study in real time [5].

Presented in this study is a high-speed functional imaging method using lattice light sheet microscopy to determine the role of lipid order and cholesterol within parasite associated membranes during invasion. Using ratio metric fluorescence imaging we show, for the first time, temporal changes in the physical properties of various membrane structures formed by the invading parasite. In addition, a computational framework to measure geometric features, such as membrane curvature has been developed. The combined spatial and temporal resolution of the lattice light sheet microscope offers unprecedented insights into the dynamic processes of *Plasmodium* invasion of red blood cells.

References

- [1] J. C. Volz, "Essential role of the PfRh5/PfRipr/CyRPA complex during *Plasmodium Falciparum* invasion of Erythrocytes," *Cell Host and Microbiome*, vol. 20, pp. 60-71, 2016.
- [2] G. E. Weiss, "Revealing the Sequence and Resulting Cellular Morphology of Receptor-Ligand Interactions during *Plasmodium falciparum* Invasion of Erythrocytes," *Plos Pathogens*, vol. 11, no. 2, 201.
- [3] D. T. Riglar, "Super-resolution dissection of coordinated events during malaria parasite invasion of the human erythrocyte," *Cell host and microbiome*, vol. 9, no. 1, pp. 9-20, 2011.
- [4] S. Dasgupta, "Membrane-wrapping contributions of the malaria parasite invasion of the human erythrocyte," *Biophysical Journal*, vol. 107, pp. 43-54, 2014.
- [5] C. Gruring, "Development and host cell modifications of *Plasmodium Falciparum* blood stages in four dimensions," *Nature Communications*, vol. 2, no. 165, 2011.