

Overexpressed LC3B and GABARAP label autophagic structures of similar size but different shape

Iman Abdollahzadeh^{1,2}, Johnny Hendriks¹, Julia Sanwald³, Indra Simons³, Oliver H. Weiergräber², Silke Hoffmann², Dieter Willbold^{2,3}, and Thomas Gensch¹

1. Institute of Complex Systems – Cellular Biophysics (ICS-4)

2. Institute of Complex Systems – Structural Biochemistry (ICS-6)

Forschungszentrum Jülich GmbH,

Wilhelm-Johnen-Straße, 52428, Jülich, Germany

3. Institut für Physikalische Biologie (IPB),

Heinrich-Heine-Universität, 40225, Düsseldorf Germany

E-mail : i.abdollahzadeh@fz-juelich.de

KEY WORDS: Atg8, GABARAP, LC3B, single molecule localization microscopy (SMLM), EYFP blinking, size distribution, shape distribution, autophagy, autophagosome biogenesis.

Autophagic structures containing mammalian autophagy-related proteins GABARAP and LC3B, two members of Atg8 family, have been investigated with conventional wide-field fluorescence and single molecule localization microscopy (SMLM), employing transient overexpression of both proteins as EYFP fusion proteins. On the basis of the well-known blinking of EYFP, super-resolution images of the corresponding GABARAP- and LC3B-containing structures with resolution of well below 50 nm were generated. These structures were quantitatively investigated (size, shape) for 10 representative cells and classified accordingly. While size distributions of structures labelled by the two proteins were found to be similar, shape distributions appeared quite disparate, with EYFP-GABARAP favouring circular structures and elliptical structures being dominant for EYFP-LC3B. The latter also featured a nearly doubled fraction of U-shaped structures. The experimental results point towards highly differential localisation of the two proteins, which appear to label structures representing distinct stages or even specific channels of autophagy and vesicular trafficking pathways. Our data also demonstrate that the application of super-resolution techniques expands the possibilities of fluorescence-based methods in autophagy studies and in some cases can rectify conclusions obtained from conventional fluorescence microscopy with diffraction-limited resolution.