

MUELLER-MATRIX OPTICAL SCANNING MICROSCOPY: FROM MICRO- TO MACROSCOPIC BIOLOGICAL ORGANIZATION

Aymeric Le Gratiet¹, Riccardo Marongiu^{1,2}, Muhammad Waseem Ashraf^{1,2}, Artemi Bendandi^{1,2}, Nicholas Anthony¹, Paolo Bianchini^{1,3}, Alberto Diaspro^{1,2}

¹ Nanoscopy, CHT Erzelli, Istituto Italiano di Tecnologia, 16152 Genova, Italy

² DIFILAB, Department of Physics, University of Genoa, Via Dodecaneso 33, 16146 Genova, Italy

³ Nikon Imaging Center, Istituto Italiano di Tecnologia, Via Morego 30, 16163 Genoa, Italy

E-mail: aymeric.legratiet@iit.it

KEY WORDS: scanning microscopy, Mueller-matrix, polarization, polarimetry, nuclear organization, zebrafish.

Microscopy techniques based on the measurements of polarimetric contrasts has proven to be an interesting tool for extracting additional information about the organization and orientation of any anisotropic sample [1]. Among these techniques, we have implemented two polarimetric multimodal optical scanning microscopy methods allowing the acquisition of all the physical effects induced by the interaction of the polarized light with a medium at different biological level of organization. This interaction can be summarized in a single 4x4 elements Mueller-matrix by comparing the independent coding and decoding polarization states, giving access simultaneously to physical parameters such as dichroism, birefringence and scattering.

First, we proposed to image sub-microscopic structures based on the measurement of the scattering differential signal from the circular left and right polarization states excitation, named Circular Intensity Differential Scattering (CIDS) [2]. Here, coupled with fluorescence imaging, we demonstrate our capability to distinguish the different chiral order conformation in chromatin-DNA from fixed isolated cell nuclei, i.e. the more compacted heterochromatin from the euchromatin areas [3].

Second, using Stokes-Mueller formalism, we demonstrate the potentiality to characterize the polarimetric transformation of non-labeled small animal. As a proof of principle, we have imaged particular biological structures of interests (collagen fibers and muscles). Then, we study the polarimetric transformation of zebrafish preserved at different embryonic developmental stages [4]. By combining the full polarimetric measurements with statistical analysis, it is possible to quantify the localized structural changes. We show that this easy and convenient method can pave the way for a better understanding of developmental biology, in which label-free techniques become a standard tool to study organisms.

References

- [1] A. Le Gratiet, et al., Optics Letters, 41(18), 2016.
- [2] A. Diaspro, et al., IEEE Trans. Biomed. Eng. 38, 670, 1991.
- [3] A. Le Gratiet, et al., OSA Continuum 1(3), 2018.
- [4] A. Le Gratiet, et al., Scientific Reports 9(1), 2019.