

NEW FLUORESCENT PROBES FOR SUPER-RESOLUTION IMAGING OF PATHOLOGICAL PROTEIN AGGREGATES IN NEURODEGENERATION.

Dr Amandeep Kaur, A/Prof. Elizabeth J. New and A/Prof Margaret Sunde

The University of Sydney, School of Medical Sciences, NSW 2006, Australia

Email: a.kaur@sydney.edu.au

KEYWORDS: Fluorescent sensors, amyloid, protein aggregation, super-resolution imaging.

In recent years, super-resolution microscopy has revolutionised the study of biological and synthetic nanostructures by breaking the diffraction limit, and allowing visualisation of cells and materials on the molecular scale.¹ While this Nobel Prize-winning technology demonstrates great potential for biomedical researchers wishing to unravel molecular-level interactions, the quantitative information obtainable and the progress in this field, is greatly limited by the limited number and poor photophysical properties of existing fluorophores.²

Protein misfolding and aggregation are hallmarks of many neurodegenerative diseases.³ A key question in this field is the nature of the differences between globular and fibrillar aggregates of these proteins and how the histopathologic patterns relate to the different types of FTLD. Therefore, an ability to visualise the molecular-level organisation, structure and distribution of proteins in these proteinopathies is cardinal to gaining deeper understanding of the mechanisms underlying the associated neurodegeneration. As will be described in the presentation, we have developed novel coumarin-based fluorescent sensors (AmyBlink) that utilise the amyloidophilic properties of the benzothiazole scaffold. AmyBlink-1 exhibits dramatic change in their emission profiles: turn-on in fluorescence and blue-shifted emission maxima. In addition, the specialised conditions that allow these sensors to under photoswitching properties crucial for super-resolution imaging will also be discussed. These sensors are highly valuable tools to researchers working towards developing improved molecular tools for super-resolution imaging and understanding neurodegenerative pathologies.




Figure 1: Differential fluorescence response of AmyBlink-1 towards diverse amyloid-forming proteins.

References

1. Patterson, G., Davidson, M., Manley, S. & Lippincott-Schwartz, J. Superresolution imaging using single-molecule localization. *Annu. Rev. Phys. Chem.* **61**, 345–367 (2010).
2. Durisic, N., Cuervo, L. L. & Lakadamyali, M. Quantitative super-resolution microscopy: pitfalls and strategies for image analysis. *Curr. Opin. Chem. Biol.* **20**, 22–28 (2014).
3. Dobson, C. M. Protein folding and misfolding. *Nature* **426**, 884 (2003).